arXiv:hep-th/9409059v1 11 Sep 1994

UMD-PP-95-43

The Inhomogeneity Expansion for Planar Q.E.D in a Magnetic Background

Gil Gat[1], Alpan Raval[2] and Rashmi Ray[3]

Physics Department
University of Maryland
College Park, MD 20742
U.S.A

Abstract

The effective action for Q.E.D in external magnetic field is constructed using the method of inhomogeneity expansion. We first treat the non-relativistic case where a Chern-Simons like term is generated. We then consider the full relativistic theory and derive the effective action for the A_μ fields. In the non-relativistic case we also add a 4-fermi type interaction and show that under certain circumstances, it corresponds to a Zeeman type term in the effective action.

[1] ggat@delphi.umd.edu
[2] raval@umdhep.umd.edu
[3] rray@delphi.umd.edu

1. **Introduction**

The behavior of a system of interacting planar fermions in an external magnetic field has many interesting features. In 2+1 dimensions such systems are of relevance for the description of the Quantum Hall Effect and high T_c superconductivity. In addition, we have the ulterior motive of adapting our technique to 3+1 dimensional situations. It is of particular (and potentially of practical) interest to study Abelian (or non-Abelian) theories involving fermions (in 3+1 dimensions) in the environment of strong magnetic fields. This is especially rewarding in a cosmological context, where strong magnetic fields may be present in the early universe, collapsing supernovae and perhaps superconducting strings. To give an idea of the magnitudes of the magnetic fields involved, we may say that those associated with compact astrophysical objects range between $O(10^4)$ T (magnetic white dwarfs) and $O(10^{10})$ T (supernovae)[1]. In between, neutron stars have associated magnetic fields of $O(10^8)$ T. It has also been suggested in some models for extragalactic gamma bursts that very strong magnetic fields $O(10^{13})$ T are involved [2]. The vacuum structure of gauge theories in a strong magnetic background, has been and continues to be a field of considerable interest [3], [4]. The minimal standard model also exhibits interesting features in the presence of a strong magnetic background, through the self-energy corrections of the neutrinos, through vacuum polarization and also through non-perturbative fermion number violation [5], [6]. It has also been observed [7],[8] that in certain planar fermionic theories with four-fermi interactions, a strong magnetic field, in conjunction with a finite density, induces breaking of the chiral symmetry and thereby generating dynamical fermion mass. We can address all these issues through minimal modifications of the techniques described below. The effective action in a magnetic background plays a pivotal role in all of them. This is one of our primary motivations in outlining our method in considerable detail.

Effective actions neatly encapsulate the contributions of quantum corrections to the classical action and thus permit a classical analysis of fully quantum problems, an advantage not to be lightly regarded. Given an effective action for a quantum theory, one can read off the various physical quantities of interest like effective masses, permeability and susceptibility with relative ease. In this paper, we wish to find the effective action of the system in terms of the fluctuating gauge fields that the planar fermions are subjected to, by tracing the fermions out of the partition function. The formal process of integrating

Grassman valued fields out yields a functional determinant, generically a nonlocal quantity. In order to obtain a Lagrangian comprising local fields and their derivatives, we have to expand the determinant within some controlled scheme. The derivative expansion method, also called the inhomogeneity expansion in the present context, is one such scheme. To implement this, we assume that the fluctuating fields coupled to the fermions vary slowly in space-time compared to the natural scales set by the background magnetic field. Namely, the typical momenta associated with these fluctuations are $\ll \frac{1}{l}$, where $l \equiv \frac{1}{\sqrt{eB}}$ is the magnetic length and the typical frequency is $\omega \ll \omega_c$, where $\omega_c m$ is the cyclotron frequency. This naturally sets a limit on contributions from the higher derivatives of the electromagnetic fields by filtering out and discarding the high frequency and the high momentum modes and thus enables us to write a local Lagrangian containing the desired number of derivatives of the field.

Our method obviously works well in the absence of gaps in the spectrum and fails precisely at the points where such gaps exist. Particularly, in this problem, where the spectrum is described by Landau levels, the method expectedly fails where the gap between successive levels is probed. In practical terms, we see, in our subsequent calculations, the emergence of spurious singularities in the form of derivatives of the delta function precisely at these points. In this work, we have adopted the somewhat pragmatic stance of disregarding these singularities and tacitly admitting the shortcomings of our method at these particular points. We however hasten to add that away from these particular points, derivative expansion is a perfectly reliable method and leads to a rather formulation of the effective action. If thermal effects are taken into account resulting in the smearing out of these discrete levels through thermal fluctuations, this method can be readily applied to even the points mentioned earlier.

Although the solution of the "free" single particle problem is known in this case, the construction of an effective action (or potential) for the interacting theory is rather involved. In particular since the theory is gauge invariant to begin with, one would like to find a method or rather an expansion, that will exhibit this gauge invariance order by order in a manifest way.

In this paper we construct such a method. We apply it to the nonrelativistic test case[1] before tackling the fully relativistic theory. The basic idea of our method is the use of guiding center coordinates which are the natural variables for this problem. Using these variables, we are able to convert the problem of calculating the fermion determinant into the problem of calculating matrix elements of known commutators and of lowering and raising operators. In addition we make several unitary transformations along the way to expose the inherent gauge invariance of the system at each stage of the approximation.

---

[1] This problem was originally considered by Ray and Sakita [9] for the non zero temperature case. Our results agree with theirs for the T=0 case (see also [14])

The organization of this paper is as follows: we start with the nonrelativistic problem, establishing the notation and describing the approximations entailed and transformations made in section 2. In section 3 we present the calculations and the results for the nonrelativistic case. In section 4 we study the effect of adding a four Fermi type interaction to the original Lagrangian and show that in a nonrelativistic theory this kind of term, under certain circumstances, can induce a Zeeman type interaction in the effective Lagrangian. In section 5 we consider the fully relativistic QED . We conclude with discussions in section 6. Useful relations are deferred to the appendices.

2. **Nonrelativistic Q.E.D**

To set the stage for the calculation we start by establishing the notation. Consider the single particle Landau problem. The dynamics of a single planar particle in an uniform magnetic field normal to the plane is described by the Landau hamiltonian:

$$H_0 = \frac{1}{2m}\mathbf{P}^2$$

where $\mathbf{A}$ may be chosen in a specific gauge $\mathbf{A} = B(-y, x)$. The wave functions of the system, $\langle x|n, X\rangle$, are labeled by two parameters; n and X. n, which is an integer is the Landau level index and X, a continuous parameter, measures the degeneracy of any given Landau level. The single particle energies are given by:

$$E_n = (n + \frac{1}{2})w_c \tag{1}$$

where $w_c = \frac{eB}{m}$ is the cyclotron frequency. These energies are obviously independent of X and the operator $\hat{X}$ which measures the value of X commutes with the hamiltonian. We call $\hat{X}$ and its conjugate $\hat{Y}$ the guiding center coordinates.They play a central role in our subsequent discussions.

Let us define $\hat{\mathbf{\Pi}}_1 = \hat{p}_1 + \frac{eb}{2}\hat{\mathbf{x}}_2$ and $\hat{\mathbf{\Pi}}_2 = \hat{p}_2 - \frac{eb}{2}\hat{\mathbf{x}}_1$ where $[\hat{\mathbf{\Pi}}_1, \hat{\mathbf{\Pi}}_2] = \frac{i}{l^2}$. We can similarly define $\hat{a} = \frac{l}{\sqrt{2}}(\hat{\mathbf{\Pi}}_1 + i\hat{\mathbf{\Pi}}_2)$. The guiding center operators are then defined as

$$\hat{X} = \hat{\mathbf{x}}_1 + l^2\hat{\mathbf{\Pi}}_2 \qquad \text{and} \qquad \hat{Y} = \hat{\mathbf{x}}_2 - l^2\hat{\mathbf{\Pi}}_1 \tag{2}$$

thus $[\hat{X}, \hat{Y}] = -il^2$. These are the two sets of cannonically conjugate operators that we need:

$$\hat{a}^\dagger\hat{a}|n, X\rangle = n|n, X\rangle \qquad \hat{X}|n, X\rangle = X|n, X\rangle$$

The normalization for X is fixed by the degeneracy of the Landau levels per unit area. Namely

$$\int dX |\langle n, X|n, X\rangle|^2 = \frac{1}{2\pi l^2} \tag{3}$$

where $\langle x|n, X\rangle = \frac{1}{\sqrt{2\pi l^3}} e^{\frac{iXy}{l^2}} u_n(\frac{x-X}{l})$ and $u_n(\frac{x-X}{l})$ are the standard harmonic oscilator wave functions.

Using an euclidean metric, the partition function for the corresponding many particle system is given by:

$$\mathcal{Z} = \int D\psi D\psi^\dagger e^{-\int d\vec{x}dt\psi^\dagger\{\Pi_\tau + \frac{1}{2m}(\mathbf{\Pi})^2 - \mu\}\psi} \tag{4}$$

where $\hat{\Pi}_\tau = -i\partial_\tau - iA_0 = \hat{p}_\tau - eA_\tau$,and $\hat{\mathbf{\Pi}} = \hat{\mathbf{p}}_\tau - e\mathbf{A}$. Integrating the fermions out we get $\mathcal{Z} = e^{-S_{eff}}$ where

$$S_{eff} = -\mathrm{Tr}\log[i\hat{\Pi}_\tau + \frac{1}{2m}\hat{\mathbf{\Pi}}^2 - \mu] \tag{5}$$

The corresponding current is

$$\langle J_\mu(x)\rangle \equiv \frac{\delta S_{eff}}{\delta A_\mu(x)}$$

where $J_\mu(x) = (ie\frac{1}{i\hat{\Pi}_\tau + \frac{1}{2m}\hat{\mathbf{\Pi}}^2 - \mu}, \{\hat{\mathbf{\Pi}}_i, \frac{1}{i\hat{\Pi}_\tau + \frac{1}{2m}\hat{\mathbf{\Pi}}^2 - \mu}\})$. We translate the bra and ket to $x = 0$, consequently the operators $\hat{x}_\mu$ in J_μ get translated to $\hat{x}_\mu + x_\mu$ and we expand all the functions of $\hat{x}_\mu + x_\mu$ around x_μ (derivative expansion).

To make the gauge invariance as manifest as possible perform a succession of unitary transformations $\Gamma = WVU$, on J_μ where

$$\begin{aligned} U(x,t) &= e^{-ie(\hat{\mathbf{x}}\cdot\mathbf{A} + \frac{1}{2!}\hat{\mathbf{x}}_i\hat{\mathbf{x}}_j\partial_i\partial_j\mathbf{A}_j + \frac{1}{3!}\hat{\mathbf{x}}_i\hat{\mathbf{x}}_j\hat{\mathbf{x}}_k\partial_i\partial_j\partial_k\mathbf{A} + \ldots)} \\ V(x,t) &= e^{-ie[\hat{\tau}A_\tau(x) + \frac{\hat{\tau}^2}{2!}\dot{A}_\tau(x) + \frac{\hat{\tau}^3}{3!}\ddot{A}_\tau + \ldots} \\ W(x,t) &= e^{\frac{i}{2l^2}\hat{\mathbf{x}}_1\hat{\mathbf{x}}_2} \end{aligned} \tag{6}$$

We notice that Γ, acting on $|x_\mu = 0\rangle$, produces just a factor of unity[2]. Thus

$$\langle J_\mu(x,t)\rangle = \langle 0|\Gamma \tilde{J}_\mu \Gamma^\dagger|0\rangle. \tag{7}$$

Since A_τ is entirely fluctuating, we refer to it as a_τ, similarly the fluctuating part of $\mathbf{A}_i$ is refered to as $\mathbf{a}_i$.

After all these transformations we obtain:

$$\begin{aligned} \langle J_\tau\rangle &= ie\langle 0|\frac{1}{i\hat{\Pi}_\tau - \mu + i\Delta_\tau + \frac{1}{2m}(\hat{\mathbf{\Pi}} + \mathbf{\Delta})^2}|0\rangle \\ \langle J_i\rangle &= \frac{e}{2m}\langle 0|\left\{\hat{\mathbf{\Pi}}_i, \frac{1}{H_0 + \frac{1}{2m}(\hat{\mathbf{\Pi}}\cdot\mathbf{\Delta} + \mathbf{\Delta}\cdot\hat{\mathbf{\Pi}} + \mathbf{\Delta}^2) + i\Delta_\tau}\right\}|0\rangle \end{aligned} \tag{8}$$

where $H_0 = i\Pi_\tau + \frac{1}{2m}\hat{\mathbf{\Pi}}^2 - \mu$,

$$H_0|n,X,w\rangle = \Gamma_n|n,X,w\rangle = [iw - \mu + (n+\frac{1}{2})w_c]|n,X,w\rangle$$

and

$$\begin{aligned} \Delta_\tau &= -e[\hat{\mathbf{x}}\cdot\partial a_\tau + \frac{1}{2}\hat{\mathbf{x}}_i\hat{\mathbf{x}}_j\partial_i\partial_j a_\tau + \hat{\mathbf{x}}_i\hat{\tau}a_i a_\tau + \ldots] \\ \mathbf{\Delta}_l &= -e[\hat{\tau}\dot{a}_l + \frac{1}{2}\hat{\tau}^2\ddot{a}_l + \hat{\tau}\hat{\mathbf{x}}\cdot\partial\dot{a}_l + \frac{1}{2}\hat{\mathbf{x}}_i\mathcal{F}_{il} + \frac{1}{3}\hat{\mathbf{x}}_i\hat{\mathbf{x}}_j\partial_i\mathcal{F}_{jl} + \ldots]. \end{aligned} \tag{9}$$

H_0 can be solved exactly (its spectrum being the L.L) and the rest is treated perturbatively in the following section. In fact,

$$H_0|n,X,w\rangle = \Gamma_n|n,X,w\rangle = [iw - \mu + (n+\frac{1}{2})w_c]|n,X,w\rangle\ . \tag{10}$$

3. **Effective Action**

The most efficient way of computing the effective action is to obtain $\langle\Pi_{\mu\nu}\rangle$ the polarization tensor, from $\langle J_\mu\rangle$ and write:

$$S_{eff} = \frac{1}{2}\int dxdy\, a_\mu(x)\langle\Pi_{\mu\nu}(x-y)\rangle a_\nu(y) \tag{11}$$

where $\langle\Pi_{\mu\nu}\rangle = \frac{\delta\langle J_\mu(x)\rangle}{\delta a_\nu(y)}$. Using the remaining rotational symmetry, the only independent components of $\langle\Pi_{\mu\nu}\rangle$ that are needed to reconstruct the effective action are $\langle\Pi_{00}\rangle$, $\langle\Pi_{01}\rangle$,

---

[2] At this point we note that the transformation V fails to be unitary in the corresponding finite temperature problem due to the compactness of the domain of definition of τ , $(0 \to \beta = 1/T)$. At $T = 0$, however, V is unitary.

$\langle \Pi_{11}\rangle$ and $\langle \Pi_{12}\rangle$. In section 2 we transformed $\langle J_\mu\rangle$ to a form that allows the use of perturbation theory. Thus for example we can expand eq.(9) to get

$$\begin{aligned}\langle J_\tau\rangle & \quad ie\langle 0|\frac{1}{i\hat{p}_\tau - \mu + i\Delta_\tau + \frac{1}{2m}(\hat{\mathbf{\Pi}} + \mathbf{\Delta})^2}|0\rangle \\ &= ie\langle 0|\frac{1}{H_0}|0\rangle - ie\langle 0|\frac{1}{H_0}\left(i\Delta_\tau + \frac{1}{2m}(\hat{\mathbf{\Pi}}\cdot\mathbf{\Delta} + \mathbf{\Delta}\cdot\hat{\mathbf{\Pi}} + \mathbf{\Delta}^2)\right)\frac{1}{H_0}|0\rangle + \ldots\end{aligned} \tag{12}$$

Now the Δ_μ generically contain $\hat{x}_\mu$. We use the definitions of $\hat{X}$ and $\hat{Y}$ from section 2 to convert the $\hat{x}_\mu$ into $\hat{X}$ and $\hat{Y}$ and $\hat{\mathbf{\Pi}}_i$. The action of $\hat{X}$ and $\hat{Y}$ on $|x_\mu\rangle = 0$ (or its complex conjugate) enables us to get rid of $\hat{X}$ and $\hat{Y}$ altogether and obtain an operator that depends only on $\hat{\mathbf{\Pi}}$. Subsequently we can introduce a complete set $\{|n, X\rangle\}$ of states and integrate over X (as the operator does not contain any further guiding center coordinates) immediately to obtain the degeneracy factor $\rho_0 = \frac{1}{2\pi l^2}$. The remaining evaluation of harmonic oscillator matrix elements is straightforward.

To clarify the foregoing statements through an example, we compute the contribution proportional to f_{12} to $\langle J_\tau\rangle$. This is given by

$$\frac{ie^2}{2m}\langle 0|\frac{1}{H_0}(\hat{\mathbf{x}}_1\hat{\mathbf{\Pi}}_2 - \hat{\mathbf{x}}_2\hat{\mathbf{\Pi}}_1)\frac{1}{H_0}|0\rangle$$

now, $\hat{\mathbf{x}}_1 = \hat{X} - l^2\hat{\mathbf{\Pi}}_2$ and $\hat{\mathbf{x}}_2 = \hat{Y} + l^2\hat{\mathbf{\Pi}}_1$, therefore

$$\langle 0|\hat{X} = l^2\langle 0|\hat{\mathbf{\Pi}}_2 \qquad\qquad \langle 0|\hat{Y} = -l^2\langle 0|\hat{\mathbf{\Pi}}_2\ .$$

Also $[\hat{\mathbf{\Pi}}_1, \frac{1}{H_0}] = -\frac{i}{ml^2}\frac{1}{H_0}\hat{\mathbf{\Pi}}_2\frac{1}{H_0}$ and similarly for $\hat{\mathbf{\Pi}}_2$, $[\hat{\mathbf{\Pi}}_2, \frac{1}{H_0}] = \frac{i}{ml^2}\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\frac{1}{H_0}$. Thus the above matrix element becomes:

$$\begin{aligned}& \frac{ie^2l^2}{2m}\langle 0|[\hat{\mathbf{\Pi}}_2, \frac{1}{H_0}]\hat{\mathbf{\Pi}}_2\frac{1}{H_0} + [\hat{\mathbf{\Pi}}_1, \frac{1}{H_0}]\hat{\mathbf{\Pi}}_1\frac{1}{H_0}|0\rangle \\ =& -\frac{e^2}{2m^2}\langle 0|\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\frac{1}{H_0}\hat{\mathbf{\Pi}}_2\frac{1}{H_0} - \frac{1}{H_0}\hat{\mathbf{\Pi}}_2\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\frac{1}{H_0}|0\rangle \\ =& -\frac{e^2}{m^2}\langle 0|\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\frac{1}{H_0}\hat{\mathbf{\Pi}}_2\frac{1}{H_0}|0\rangle \\ =& -\frac{e^2}{m^2}\sum_{n_1,n_2,n_3}\int dX\int\frac{dw}{2\pi}\frac{\langle 0|n_1, X, w\rangle\langle n_3, X, w|0\rangle}{\Gamma_{n_1}\Gamma_{n_2}\Gamma_{n_3}}\langle n_1|\hat{\mathbf{\Pi}}_1|n_2\rangle\langle n_2|\hat{\mathbf{\Pi}}_2|n_3\rangle\end{aligned} \tag{13}$$

The integration over X can now be done trivially yielding $\frac{1}{2\pi l^2}\delta_{n_1,n_3}$, reducing the matrix element to:

$$-\frac{e^2}{2\pi m^2 l^2}\sum_{n_1,n_2}\int\frac{dw}{2\pi}\frac{\langle n_1|\hat{\mathbf{\Pi}}_1|n_2\rangle\langle n_2|\hat{\mathbf{\Pi}}_2|n_3\rangle}{\Gamma^2_{n_1}\Gamma_{n_2}}$$

Using the definition of $\hat{\mathbf{\Pi}}$ we get

$$-\frac{ie^2}{4\pi m^2 l^4}\int\frac{dw}{2\pi}\sum_{n=0}^{\infty}(n+1)\left[\frac{1}{\Gamma_n^2\Gamma_{n+1}}-\frac{1}{\Gamma_n\Gamma_{n+1}^2}\right] \tag{14}$$

and we use $\frac{1}{\Gamma_n\Gamma_{n+1}}=\frac{1}{w_c}\left[\frac{1}{\Gamma_n}-\frac{1}{\Gamma_{n+1}}\right]$ Thus the above matrix element reduces to (retaining only the non-singular part as discussed in the introduction)

$$\frac{ie^2}{2\pi}\sum_{n=0}^{\infty}\int\frac{dw}{2\pi}\frac{1}{iw-\mu+(n+\frac{1}{2})w_c}f_{12}(\hat{\mathbf{x}},\tau)=\frac{ie^2}{2\pi}\sum_{n=0}^{\infty}\Theta(w_c(n+\frac{1}{2})-\mu)f_{12}(\hat{\mathbf{x}},\tau)$$

For operators involving $\hat{\tau}$ we use $[\hat{\tau},\frac{1}{H_0}]=\frac{1}{H_0^2}$ and proceed as above. Another simplifying feature is that any matrix element with an odd number of $\hat{\mathbf{\Pi}}_i$ is zero.

It is easy to show that all terms which could possibly lead to gauge noninvariance vanish identically. We append an example here. The term proportional to $\ddot{a}_i$ in $\langle J_\tau\rangle$ leads to a gauge noninvariant term in the effective action. The matrix element proportional to this term is

$$\frac{ie^2}{2m}\langle 0|\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\hat{\tau}^2\frac{1}{H_0}|0\rangle\ddot{a}_1+(1\to 2) \tag{15}$$

Now since $\hat{\tau}^2\frac{1}{H_0}|0\rangle=\frac{2}{H_0^3}|0\rangle$ we get

$$\frac{ie^2}{m}\langle 0|\frac{1}{H_0}(\hat{\mathbf{\Pi}}_1\ddot{a}_1+\hat{\mathbf{\Pi}}_2\ddot{a}_2)\frac{1}{H_0}|0\rangle=0 \tag{16}$$

so that in fact this term does not appear.

At this point we collect all the relevant terms, i.e. those proportional to $\partial_1^2 a_\tau$, $\partial_1\partial_\tau a_1$ and f_{12} for $\langle J_\tau\rangle$ and proportional to $\partial_\tau^2 a_1$, $\partial_\tau^2 a_2$, $\partial_1\partial_\tau a_\tau$, $\partial_2\partial_\tau a_\tau$ and $\partial_2 f_{12}$ for $\langle J_1\rangle$ and obtain :

$$\begin{array}{rcl}
\langle\Pi_{\tau\tau}(x,y)\rangle &=& \frac{e^2}{2\pi w_c}C(\partial_1^2+\partial_2^2)\delta(x-y)+\ldots \\
\langle\Pi_{\tau 1}(x,y)\rangle &=& -\frac{e^2}{2\pi w_c}C\partial_\tau\partial_1\delta(x-y)-\frac{ie^2}{2\pi}C\partial_2\delta(x-y)+\ldots \\
\langle\Pi_{\tau 2}(x,y)\rangle &=& -\frac{e^2}{2\pi w_c}C\partial_\tau\partial_2+\frac{ie^2}{2\pi}C\partial_1\delta(x-y)+\ldots \\
\langle\Pi_{11}(x,y)\rangle &=& \frac{e^2}{2\pi w_c}C\partial_\tau^2\delta(x-y)-\frac{e^2}{\pi m}C_1\partial_2^2\delta(x-y)+\ldots \\
\langle\Pi_{12}(x,y)\rangle &=& \frac{e^2}{\pi m}C_1\partial_1\partial_2\delta(x-y)-\frac{ie^2}{2\pi}C\partial_\tau\delta(x-y)+\ldots \\
\langle\Pi_{22}(x,y)\rangle &=& \frac{e^2}{2\pi w_c}C\partial_\tau^2\delta(x-y)-\frac{e^2}{\pi m}C_1\partial_1^2\delta(x-y)+\ldots
\end{array} \tag{17}$$

where

$$C = \sum_{n=0}^{\infty} \int \frac{dw}{2\pi} \left(1 + w_c \frac{\partial}{\partial w_c} \right) \frac{1}{\Gamma_n} = \sum_{n=0}^{\infty} \Theta(\mu - (n + \frac{1}{2}) w_c) + \dots \tag{18}$$

and C_1 is

$$C_1 = \sum_{n=0}^{\infty} (n + \frac{1}{2}) \int \frac{dw}{2\pi} \left(1 + \frac{1}{2} w_c \frac{\partial}{\partial w_c} \right) \frac{1}{\Gamma_n} = \sum_{n=0}^{\infty} (n + \frac{1}{2}) \Theta(\mu - (n + \frac{1}{2}) w_c) + \dots\ . \tag{19}$$

The effective action is reconstructed as :

$$S_{eff} = \frac{ie^2}{2\pi} C \epsilon_{\mu\nu\rho} \int d^3x a_\mu \partial_\nu a_\rho + \frac{e^2}{4\pi w_c} C \int d^3x \, \mathcal{E}^2 + \frac{e^2}{2\pi m} C_1 \int d^3x \, \mathcal{B}^2 + \dots \tag{20}$$

where $\vec{\mathcal{E}}$ and $\mathcal{B}$ are the euclidean fluctuating electric and magnetic field strength respectively. If we choose μ such that exactly N Landau Levels are filled $C \to N + 1$ and $C_1 \to \frac{(n+\frac{1}{2})^2}{2}$.

We note two points regarding this effective action. The first is the emergence of a C-S term. The second is that since relativistic invariance is broken by the external magnetic field, we get different coefficients for the fluctuating $\mathcal{E}^2$ and $\mathcal{B}^2$ terms. Further if the magnetic field is very strong (i.e. $l \ll \frac{1}{m}$), the second term in the effective action does not contribute. This corresponds to dimensional reduction. We also observe that if there are no fermions ,i.e. $\mu = 0$, the magnetic $\mathcal{B}^2$ term in the gauge field action is not renormalized by quantum effects. This is to be contrasted with the relativistic case (see eq.(63)) where the vacuum (filled Dirac sea) does contribute to the renormalization of this term.

We can use this effective action (rotated to minkowski space) to obtain physically interesting quantities, for example the dielectric function

$$\epsilon(\mathbf{p}) = 1 + \frac{\Pi_{00}(0, \mathbf{p})}{\mathbf{p^2}} = 1 + \frac{e^2}{2\pi w_c} C$$

and the magnetic permeability

$$\frac{1}{\mu(0, \mathbf{p})} = 1 + \frac{\Pi_i^i(0, \mathbf{p})}{\mathbf{p^2}} = 1 - \frac{e^2}{\pi m} C_1 \ .$$

4. **Effective Zeeman Interaction**

Having obtained the effective action we take this oportunity to explore an interesting phenomenon namely the effect of adding a four fermi term. In this section we show that if a four fermi term is added to the original Lagrangian introduced in the previous sections, an effective Zeeman-type interaction could be induced in certain cases.

The partition function for the system is

$$\begin{aligned} \mathcal{Z} &= \int D\phi D\psi D\psi^\dagger e^{S(\phi,\psi\bar{\psi})} \\ S &= i\int d^3x \left[\psi^\dagger[i\partial_0 - eA_0 + \mu + \frac{1}{2m}(i\nabla + e\mathbf{A})^2]\psi - g(\psi^\dagger\frac{\sigma_3}{2}\psi)^2\right] \end{aligned} \tag{21}$$

Using an auxiliary field ϕ to disentangle the quartic term, we obtain

$$\mathcal{Z} = \int D\psi D\psi^\dagger exp\left\{i\int d^3x\left[\psi^\dagger[i\partial_0 - eA_0 + \mu + \frac{1}{2m}(i\nabla + e\mathbf{A})^2 + \sqrt{\frac{g}{2}}\phi\sigma_3]\psi + \frac{1}{2}\phi^2\right]\right\} \tag{22}$$

We wish to show that the effective potential for ϕ obtained upon integrating the fermions out is minimized by a non-zero constant configuration which we call ϕ_0. This in turn would mean that the Yukawa term in the action above yields an effective Zeeman term $\sqrt{\frac{g}{2}}\phi_0\psi^\dagger\sigma_3\psi$.

The effective potential for ϕ is given by

$$V_{eff} = -\frac{1}{2\pi l^2}\sum_{n=0}^{\infty}\int\frac{dk_0}{2\pi i}\log\left[\frac{k_0 - E_n + \mu + \frac{b}{2}}{k_0 - E_n + \mu - \frac{b}{2}}\right], \tag{23}$$

where $E_n = \frac{1}{ml^2}(n + \frac{1}{2})$ and $b \equiv \sqrt{2g}\phi$. The presence of the sum of two logarithms is due to the two component nature of the fermion. The mean density is computed from V_{eff} as $\langle 0|\rho|0\rangle = -\frac{\partial V}{\partial\mu}$, which yields

$$\langle 0|\rho|0\rangle = \frac{1}{2\pi l^2}\sum_{n=0}^{\infty}\int\frac{dk_0}{2\pi i}\left[\frac{1}{k_0 - E_n + \mu + \frac{b}{2}} + \frac{1}{k_0 - E_n + \mu - \frac{b}{2}}\right]. \tag{24}$$

On performing the k_0 integral, we get

$$\langle 0|\rho|0\rangle \equiv \langle 0|\rho|0\rangle_+ + \langle 0|\rho|0\rangle_-, \tag{25}$$

where

$$\langle 0|\rho|0\rangle_+ \equiv \frac{1}{2\pi l^2}\sum_{n=0}^{\infty}\theta(\mu + \frac{b}{2} - E_n) \tag{26}$$

$$\langle 0|\rho|0\rangle_- \equiv \frac{1}{2\pi l^2}\sum_{n=0}^{\infty}\theta(\mu - \frac{b}{2} - E_n) \tag{27}$$



Figure 1: Graphical solution of eq.(30). The y axis is labeled in units of $\rho_0 = \frac{eB}{2\pi}$

We reconstruct V_{eff} from $\langle 0|\rho|0\rangle$ using

$$V_{eff} = -\langle 0|\rho|0\rangle A_0 - \frac{1}{4g}b^2, \tag{28}$$

where the second term is the tree level contribution. We note that in the absence of fluctuating electrmagnetic fields, the role of A_0 is played by $-\frac{1}{2e}b$ in $\langle 0|\rho|0\rangle_+$ and by $\frac{1}{2e}b$ in $\langle 0|\rho|0\rangle_-$. Thus

$$V_{eff}(b) = \frac{1}{2e}b\left(\langle 0|\rho|0\rangle_+ - \langle 0|\rho|0\rangle_-\right) - \frac{1}{4g}b^2. \tag{29}$$

We wish to look for extrema of the above away from the points $b = \pm[(2n+1)\frac{1}{ml^2} - 2\mu]$, since at these points the extrema picks up singular delta function contributions. The extrema satisfy

$$\langle 0|\rho|0\rangle_+ - \langle 0|\rho|0\rangle_- = \frac{e}{g}b. \tag{30}$$

Since the l.h.s. of eq.(30) is a complicated function of b, it is solved graphically (see fig.(1)). It is seen that if the external magnetic field is strong enough to satisfy $B^2 > \frac{m^2 g}{\pi e^3}$, at least one non-zero solution exists which minimizes the effective potential. One such solution, which is for the situation where $B^2 \simeq \frac{m^2 g}{\pi e^3}$, is $b_0 = \omega_c$. This, as has been discussed earlier in this section, leads to an effective Zeeman interaction term $\frac{\omega_c}{2}\psi^\dagger\sigma_3\psi$.

Thus, we have the rather interesting result that a four fermi term in a strong external magnetic field can lead to an effective Zeeman interaction for the fermions.

5. **Relativistic Q.E.D.**

Having demonstrated the efficacy of our technique for the nonrelativistic case, we now go on to discuss the relativistic scenario. Although this is deemed unnecessary for most condensed matter oriented applications, there are situations where the fermionic dispersion relation may be taken to be approximately linear (e.g. in the neighbourhood of the Fermi surface) and our subsequent calculations would find applicability in precisely these cases. Furthermore, as has been discussed at length in the introduction, the astrophysical and cosmological ramifications of this problem, namely fermions interacting through a gauge field, Abelian or otherwise, in a strong magnetic background, are manifold. The strong magnetic field can induce fundamental changes not only in the vacuum structure of such theories but also in the other sectors of the Fock space.

Admittedly, in this work, we have addressed the problem of only planar fermions with an Abelian interaction. However, adding on the third spatial dimension entails only very minor modifications, namely, the single particle energies also come to depend quadratically on the z-component of the momentum. This can be incorporated quite easily within our computational framework as can the extension to non-Abelian interactions, which actually bring in little more than notational complexity.

Having enumerated these motivations, we now proceed to the actual computation. The generating functional is then given by

$$\mathcal{Z} = \int D\psi D\bar{\psi} e^{-\int d\mathbf{x} dt \bar{\psi}\{\not{D} - m - \mu\gamma^0\}\psi} \equiv e^{-S_{eff}} \tag{31}$$

where μ is the chemical potential and we use an euclidean metric with signature -1. The fermionic degrees of freedom can then be integrated out to yield the formal expression of the effective action as

$$S_{eff} = -\mathrm{Trlog}[\not{D} - m - \mu\gamma^0] \tag{32}$$

Define $\tilde{a}_0 = a_0 - \frac{i\mu}{e}$, and $\tilde{\mathbf{a}} = \mathbf{a}$. Then we can express the effective action as

$$S_{eff} = -\mathrm{Trlog}[\tilde{\not{D}} - m] \,, \tag{33}$$

with

$$\tilde{D}_\mu = ip_\mu - ie\tilde{a}_\mu \equiv i\Pi_\mu . \tag{34}$$

Note that the commutator $[\Pi_1, \Pi_2] = \frac{i}{l^2}$.

Upon functionally differentiating with respect to the gauge fields, the corresponding current is obtained as

$$\langle J_\mu \rangle = \frac{ie}{2} \mathrm{Tr} \langle x | \{ \gamma_\mu , \frac{1}{i \not\Pi - m} \} | x \rangle \tag{35}$$

The derivative expansion can now be performed in an identical manner to the nonrelativistic case. The bra and ket are translated to the spacetime origin, and the operators $\hat{x}_\mu$ to $\hat{x}_\mu + x_\mu$. We then formally expand all functions of $\hat{x}_\mu + x_\mu$ around x_μ. Also, the unitary transformations U, V, and W on the current operator are identical to those in the non-relativistic case. Upon performing these transformations the current takes the form:

$$\langle J_\mu \rangle = \frac{ie}{2} Tr \langle 0 | \{ \gamma_\mu , \frac{1}{i \not\Pi - m - \not O} \} | 0 \rangle \tag{36}$$

with

$$O^0 = ie \hat{x}_i \partial^i a^0 + \frac{ie}{2} \hat{x}_i \hat{x}_j \partial^i \partial^j a^0 + ie \hat{x}_0 \hat{x}_i \partial^i \partial^0 a^0 + \ldots \tag{37}$$

and

$$O^l = \frac{ie}{2} \hat{x}_i \partial^i a^l + \frac{ie}{3} \hat{x}_i \hat{x}_j \partial^i \partial^j a^l + ie \hat{x}_0 \partial^0 a^l + ie \hat{x}_0 \hat{x}_i \partial^0 \partial^i a^l + \frac{ie}{2} \hat{x}_0^2 (\partial^0)^2 a^l \tag{38}$$

Again, we can use perturbation theory to expand :

$$\frac{1}{i \not\Pi - m - \not O} = \frac{1}{i \not\Pi - m} + \frac{1}{i \not\Pi - m} \not O \frac{1}{i \not\Pi - m} + \ldots \tag{39}$$

Also, the algebraic relations between γ matrices (see appendix **A**) can be used to express:

$$\frac{1}{i \not\Pi - m} = \frac{1}{\Pi^2 - m^2 + \frac{1}{l^2} \gamma^0} (i \not\Pi + m) \tag{40}$$

Defining

$$\begin{array}{rcl} S & = & \Pi^2 - m^2 \\ M_0 & = & (\Pi^2 - m^2)^2 - \frac{1}{l^4} \\ Q_1 & = & \frac{S}{M_0} \\ Q_2 & = & \frac{1}{l^2 M_0} \end{array} \tag{41}$$

we obtain

$$\frac{1}{i\not{\Pi} - m} = (Q_1 + \gamma_0 Q_2)(i\not{\Pi} + m) \tag{42}$$

and therefore

$$\begin{aligned} \langle J_\mu(x) \rangle &= \frac{ie}{2} Tr \langle 0 | \{ \gamma_\mu, (Q_1 + \gamma_0 Q_2)(i\not{\Pi} + m) \\ &+ (Q_1 + \gamma_0 Q_2)(i\not{\Pi} + m) \not{O} (Q_1 + \gamma_0 Q_2)(i\not{\Pi} + m) \} | 0 \rangle + \ldots \end{aligned} \tag{43}$$

In order to calculate expectation values in the state $|0\rangle$, it will be necessary to expand this state in the eigenbasis of the operator Π^2. It is instructive to compute the eigenspectrum of Π^2 explicitly since this calculation will also motivate the use of guiding center coordinates .

Consider the eigenvalue equation

$$(\Pi^2 - m^2)\Psi = \lambda \Psi \tag{44}$$

i.e.

$$[(p_0 + i\mu)^2 + p_1^2 + (p_2 - eBx_1)^2]\Psi = -(\lambda + m^2)\Psi \tag{45}$$

One can solve this by separation of variables:

$$\Psi(x_1, x_2, x_0) = e^{-i\omega x^0} e^{-ieBXx^2} \Phi(x_1) \tag{46}$$

This leads to the single differential equation for Φ:

$$[-\partial_1^2 + (eB)^2 (x^1 - X)^2]\Phi = -[\lambda + m^2 + (\omega - i\mu)^2]\Phi \tag{47}$$

This is just the Schrödinger equation for a harmonic oscillator in one dimension., with eigenvalue spectrum $(2n+1)/l^2$. The full eigenvalue spectrum of the operator $\Pi^2 - m^2$ is therefore

$$\lambda_n = -\frac{(2n+1)}{l^2} - m^2 - (\omega - i\mu)^2 \tag{48}$$

The corresponding eigenstates may be labelled by the quantum numbers n, ω and X. It follows that

$$\begin{aligned} S|X, n, \omega\rangle &= \lambda_n |X, n, \omega\rangle & (49) \\ M_0 |X, n, \omega\rangle &= \Gamma_n |X, n, \omega\rangle & (50) \end{aligned}$$

where $\Gamma_n = \lambda_n^2 - \frac{1}{l^4}$ Therefore,

$$\begin{aligned} Q_1|X,n,\omega\rangle &= \frac{\lambda_n}{\Gamma_n}|X,n,\omega\rangle \\ Q_2|X,n,\omega\rangle &= \frac{1}{l^2\Gamma_n}|X,n,\omega\rangle \end{aligned} \tag{51}$$

and the completeness relation takes the form

$$\sum_n \int \frac{d\omega}{2\pi} dX|X,n,\omega\rangle\langle X,n,\omega| = 1 \tag{52}$$

We now define the operators

$$\begin{aligned} \hat{X} &\equiv -\hat{x}_1 - l^2\Pi_2 \equiv -l^2 p_2 \\ \hat{Y} &\equiv -\hat{x}_2 + l^2\Pi_1 \equiv -\hat{x}_2 + l^2 p_1 \end{aligned} \tag{53}$$

They have the properties

$$\begin{aligned} \hat{X}|X,n,\omega\rangle &= X|X,n,\omega\rangle \\ [\hat{X},\hat{Y}] &= il^2 \\ \hat{X}|0\rangle &= -l^2\Pi_2|0\rangle \\ \hat{Y}|0\rangle &= l^2\Pi_1|0\rangle \end{aligned} \tag{54}$$

As we shall later see, these properties give us a powerful method of computing expectation values of quantities in the state $|0\rangle$.

Consider eq.(44). Since the trace operation is independent of the ordering of operators in the commutator, the expression for the current to first order in O is given by:

$$\langle J_\mu\rangle = ie\mathrm{Tr}\langle 0|\gamma_\mu(Q_1+\gamma_0 Q_2)(i\not{\Pi}+m)\not{O}(Q_1+\gamma_0 Q_2)(i\not{\Pi}+m)|0\rangle \tag{55}$$

This gives rise to 16 separate terms. The calculation is done in the following manner: First, in each term, the trace is taken with respect to the γ matrices. This leaves products of Q's ,Π's and O in each term. The dependence on the x_μ in O is removed through the use of properties eq.(54) of the guiding center coordinates, as will be shown in a specific example below. Then the state $|0\rangle$ is expanded in the eigenstates of Q_1. Remembering that $\Pi_{1,2}$ act as combinations of raising and lowering operators on these eigenstates, the expectation value can be computed in a straightforward, albeit tedious, manner.

To illustrate, consider the term $-ie\langle 0|\gamma_0 Q_1 \not{\Pi}\hat{O}\gamma_0 Q_2 \not{\Pi}|0\rangle$ in $\langle J_0\rangle$. Upon taking the trace with respect to the γ matrices, this splits into 6 separate terms:

$$\begin{aligned} &2e\langle 0|Q_1\Pi_1 O^0 Q_2\Pi_2|0\rangle - 2e\langle 0|Q_1\Pi_2 O^0 Q_2\Pi_1|0\rangle - 2e\langle 0|Q_1\Pi_0 O^1 Q_2\Pi_2|0\rangle \\ &+2e\langle 0|Q_1\Pi_0 O^2 Q_2\Pi_1|0\rangle + 2e\langle 0|Q_1\Pi_1 O^2 Q_2\Pi_0|0\rangle - 2e\langle 0|Q_1\Pi_2 O^1 Q_2\Pi_0|0\rangle \end{aligned} \tag{56}$$

We would like to obtain a contribution proportional to f^{12} from these terms as a demonstration. We note that contributions to f^{12} are obtained from O_1 and O_2:

$$O_1 \rightarrow \frac{ie}{2}\hat{\mathbf{x}}_2 f^{12} \qquad\qquad O_2 \rightarrow -\frac{ie}{2}\hat{\mathbf{x}}_1 f^{12}$$

Thus the terms from eq.(56) which contribute to f^{12} are:

$$ie^2[\langle 0|Q_1\hat{\mathbf{x}}_2 Q_2\hat{\mathbf{\Pi}}_2\hat{\Pi}_0|0\rangle + \langle 0|Q_1\hat{\mathbf{\Pi}}_2\hat{\mathbf{x}}_2 Q_2\hat{\Pi}_0|0\rangle + \langle 0|Q_1\hat{\mathbf{x}}_1 Q_2\hat{\mathbf{\Pi}}_1\hat{\Pi}_0|0\rangle + \langle 0|Q_1\hat{\mathbf{\Pi}}_1\hat{\mathbf{x}}_1 Q_2\hat{\Pi}_0|0\rangle]f^{12}$$

Using eq.(54) and the commutators in appendix **B** we get

$$-\frac{4e^2}{l^2}[\langle 0|\frac{S}{M_0}\{S,\hat{\mathbf{\Pi}}_1\}\frac{1}{M_0^2}\hat{\mathbf{\Pi}}_1\hat{\Pi}_0|0\rangle - \langle 0|\frac{S}{M_0^2}\hat{\mathbf{\Pi}}_1\frac{1}{M_0}\{S,\hat{\mathbf{\Pi}}_1\}\hat{\Pi}_0|0\rangle - \langle 0|\hat{\mathbf{\Pi}}_1\frac{1}{M_0^2}\hat{\mathbf{\Pi}}_1\hat{\Pi}_0|0\rangle]f^{12}.$$

Now there is another term in $\langle J_0\rangle$ with $Q_1 \rightarrow Q_2$. This contributes to f^{12} in the form:

$$-\frac{4e^2}{l^2}[\langle 0|\frac{1}{M_0}\{S,\hat{\mathbf{\Pi}}_1\}\frac{S}{M_0^2}\hat{\mathbf{\Pi}}_1\hat{\Pi}_0|0\rangle + \langle 0|\frac{1}{M_0^2}\hat{\mathbf{\Pi}}_1\frac{S}{M_0}\{S,\hat{\mathbf{\Pi}}_1\}\hat{\Pi}_0|0\rangle - \langle 0|\frac{1}{M_0}\hat{\mathbf{\Pi}}_1^2\frac{1}{M_0}\hat{\Pi}_0|0\rangle]f^{12}.$$

Consider the simplest term:

$$\begin{aligned} &\frac{4e^2}{l^2}f^{12}\langle 0|\hat{\mathbf{\Pi}}_1\frac{1}{M_0^2}\hat{\mathbf{\Pi}}_1\hat{\Pi}_0|0\rangle = \\ &\frac{4e^2}{l^2}f^{12}\sum_{n,n_1,n_2}\int\frac{dw}{2\pi}(w-i\mu)\int dX\frac{\langle 0|n,X,w\rangle\langle n|\hat{\mathbf{\Pi}}_1|n_1\rangle\langle n_1|\hat{\mathbf{\Pi}}_1|n_2\rangle\langle n_2,X,w|0\rangle}{\Gamma_{n_1}^2} \end{aligned} \tag{57}$$

The integration over X now gives ρ_0 so we get

$$\begin{aligned} \frac{2e^2}{\pi l^4}f^{12}\sum_{n_1,n} &\int \frac{dw}{2\pi}(w-i\mu)\frac{\langle n|\hat{\mathbf{\Pi}}_1|n_1\rangle\langle n_1|\hat{\mathbf{\Pi}}_1|n\rangle}{\Gamma_{n_1}^2} \\ &= \frac{e^2}{\pi l^6}f^{12}\sum_n\int\frac{dw}{2\pi}(w-i\mu)\left[\frac{n+1}{\Gamma_{n+1}^2}+\frac{n}{\Gamma_{n-1}^2}\right] \\ &= \frac{e^2}{\pi l^6}f^{12}\sum_n\int\frac{dw}{2\pi}(w-i\mu)(n+1)\left[\frac{1}{\Gamma_{n+1}^2}+\frac{1}{\Gamma_n^2}\right] \end{aligned} \tag{58}$$

simplifying the other terms in the same way we get

$$-\frac{2e^2}{\pi l^6} f^{12} \sum_{n=0}^{\infty} \int \frac{dw}{2\pi} (w - i\mu) \frac{(n+1)(\lambda_n + \lambda_{n+1})}{\Gamma_n \Gamma_{n+1}} \left(\frac{\lambda_n}{\Gamma_n} + \frac{\lambda_{n+1}}{\Gamma_{n+1}} \right)$$

The coefficient of the term proportional to f^{12} in $\langle j_0 \rangle$ also receives contributions proportional to m. These are calculated to be

$$\frac{2ie^2}{\pi l^2 w_c} f^{12} \sum_{n=0}^{\infty} \int \frac{dw}{2\pi} \left[(n+1) \frac{(\lambda_n + \lambda_{n+1})}{\Gamma_n \Gamma_{n+1}} \left(\frac{\lambda_n (\lambda_{n+1} - \frac{1}{l^2})}{\Gamma_n} + \frac{\lambda_{n+1}(\lambda_n - \frac{1}{l^2})}{\Gamma_{n+1}} \right) - (2n+1) \frac{\lambda_n}{\Gamma_n^2} \right] \tag{59}$$

A further simplification is in order in the above expressions. Let us define

$$d_n \equiv -\frac{2n}{l^2} - m^2 - (w - i\mu)^2 \,.$$

Thus

$$\frac{1}{\Gamma_n} = \frac{1}{\lambda_n^2 - \frac{1}{l^4}} = \frac{l^2}{2} \left(\frac{1}{d_{n+1}} - \frac{1}{d_n} \right) \tag{60}$$

using this technique, any denominator containing arbitrary products of the Γs can be simplified to powers of a given d. Further, the numerators can trivially rewritten in terms of d since $\lambda_n = d_n - \frac{1}{l^2}$.

The final expression for $\langle J_0 \rangle$ and $\langle J_1 \rangle$ can be expressed entirely in terms of w , μ and d_n. It is quite straightforward to show, as in the nonrelativistic case, that the terms that potentially violate gauge invariance vanish identically. Compiling all the terms together we get:

$$\begin{aligned} \langle J_0 \rangle &= C_0 f^{12} + C_1 \partial^0 \partial^1 a^1 + C_2 (\partial^1)^2 a^0 + \ldots \\ \langle J_1 \rangle &= D_0 \partial^2 f^{12} + D_1 \partial^0 \partial^1 a^0 + D_2 (\partial^0)^2 a^1 + \ldots \end{aligned} \tag{61}$$

where C_0 , C_i and D_0 , D_i are constants to be specified later. Correspondingly,

$$\begin{aligned} \langle \Pi_{00}(x,y) \rangle &= C_2 (\partial^1)^2 \delta(x-y) + C_2 (\partial^2)^2 \delta(x-y) + \ldots \\ \langle \Pi_{01}(x,y) \rangle &= -C_0 \partial^2 \delta(x-y) + C_1 \partial^0 \partial^1 \delta(x-y) + \ldots \\ \langle \Pi_{11}(x,y) \rangle &= -C_0 (\partial^2)^2 \delta(x-y) + D_2 (\partial^0)^2 \delta(x-y) + \ldots \\ \langle \Pi_{12}(x,y) \rangle &= -D_0 \partial^2 \partial^1 \delta(x-y) + \ldots \\ \langle \Pi_{02}(x,y) \rangle &= C_0 \partial^1 \delta(x-y) + C_1 \partial^0 \partial^2 \delta(x-y) + \ldots \\ \langle \Pi_{22}(x,y) \rangle &= -D_0 (\partial^1)^2 \delta(x-y) + D_2 (\partial^0)^2 \delta(x-y) + \ldots \end{aligned} \tag{62}$$

Gauge invariance forces $C_1 = -C_2 = D_1 = -D_2 \equiv C$ so that the effective action that one gets after collecting all the terms and functionally integrating $\langle \Pi_{\mu\nu} \rangle$ is

$$S_{eff} = \frac{C}{2} \int d^3x \, \mathcal{E}^2 + \frac{D}{2} \int d^3x \, \mathcal{B}^2 + \frac{C_0}{2} \int d^3x \, a^0 \mathcal{B} + \ldots \tag{63}$$

where

$$D_0 \equiv D = -\frac{e^2}{12\pi} \sum_{n=0}^{\infty} \left[\frac{m}{2w_c} + 7n - \frac{5}{2} \right] \frac{1}{\sqrt{m^2 + 2neB}} \Theta(\sqrt{m^2 + 2neB} - |\mu|) \ , \tag{64}$$

$$C_0 = \frac{e^2}{2\pi} \sum_n \int dw \, \frac{w - i\mu}{d_n} = -\frac{ie^2}{2} \frac{\mu}{|\mu|} \sum_n \Theta(|\mu| - \sqrt{2meB + m^2})$$

and

$$\begin{aligned} C \quad &= \quad \frac{e^2 m}{32\pi w_c} \sum_{n=0}^{\infty} \frac{(10 - 13n)}{\sqrt{m^2 + 2neB}} \Theta(\sqrt{m^2 + 2neB} - |\mu|) \\ &- \quad \frac{e^2}{16\pi} \sum_{n=0}^{\infty} \frac{n(10 - 13n)}{\sqrt{m^2 + 2neB}} \Theta(\sqrt{m^2 + 2neB} - |\mu|) \end{aligned} \tag{65}$$

One sees that even in the relativistic case the coefficient of the euclidean $\mathcal{E}^2$ vanishes for strong external magnetic field. The explicit dependence on the chemical potential μ can be removed by specifying the number of filled Landau levels. Physical quantities can be extracted from this action as in the nonrelativistic case.

Discussions

In this paper we have calculated the effective action of nonrelativistic and relativistic 2+1 dimensional QED in a strong external magnetic field using the inhomogeneity expansion method. We have explicitly addressed only the case of the parity odd model i.e. one fermion flavour , but it is straightforward to use our method in the parity even case by considering two fermion flavours with positive and negative mass [15]. For the nonrelativistic case we obtain a Chern-Simons like term in addition to the renormalization of the coefficients of $\mathcal{E}^2$ and $\mathcal{B}^2$. In the relativistic case a true Chern-Simons term is generated and we calculate its coefficient as well as the leading term in the coefficients of $\mathcal{E}^2$ and $\mathcal{B}^2$. For very stong external magnetic fields ($l \ll \frac{1}{m^2}$) we see that the coefficient of the euclidean $\mathcal{E}^2$ term vanishes both in the relativistic and nonrelativistic case. This suggests that a dimensional reduction is indeed taking place [12]. In this context, it would be interesting to study the effect of such a reduction in the case of Nambu Jona-Lasinio models in general and the 2+1 dimensional Thirring model in particular [13].

We have extracted just a few of the many possible physical quantities that can be obtained from these effective actions *viz* the dielectric constant and magnetic permeability. Considering a finite temperature variant [14], we can further discuss the physics of plasmons in a magnetic field background. A particularly intriguing direction is to adapt these techniques to astrophysical and cosmological applications.

Acknowledgements

We wish to thank Jonathan Simon for interesting discussions. This work is partially supported by the N.S.F.

Appendix A. Dirac algebra and Trace relations

In the relativistic calculation, the metric we use is $g_{\mu\nu} = \mathrm{diag}(-1,-1,-1)$. One has to be careful, therefore, in raising and lowering indices. Every raising or lowering of an index picks up a negative sign. The canonical commutation relation takes the form

$$[\hat{x}_\mu, \hat{p}_\nu] = ig_{\mu\nu} \tag{66}$$

from which commutation relations involving Π's follow directly.

The algebra of Dirac matrices in 2+1 dimensions, with a negative definite Euclidean metric, albeit straightforward, is not generally familiar. Here we quote the main relations which are used in the body of the paper.

To start with, the Dirac matrices are defined as $\gamma_{1,2} = \sigma_{1,2}$, $\gamma_0 = \sigma_3$, where the σ's are the usual Pauli matrices. (note that $\gamma^i = -\sigma_i$, because the metric has negative definite signature.) This leads to the relations

$$\gamma_\mu\gamma_\nu = -g_{\mu\nu} + i\epsilon_{\mu\nu\rho}\gamma_\rho \tag{67}$$

Hence

$$\{\gamma_\mu, \gamma_\nu\} = -2g_{\mu\nu}. \tag{68}$$

The trace relations are obtained as

$$\begin{aligned} \mathrm{Tr}\gamma_\mu\gamma_\nu &= -2g_{\mu\nu} \\ \mathrm{Tr}\gamma_\mu\gamma_\nu\gamma_\lambda &= 2i\epsilon_{\mu\nu\lambda} \\ \mathrm{Tr}\gamma_\mu\gamma_\nu\gamma_\alpha\gamma_\beta &= 2(g_{\mu\nu}g_{\alpha\beta} - g_{\mu\alpha}g_{\nu\beta} + g_{\mu\beta}g_{\nu\alpha}) \end{aligned} \tag{69}$$

$$
\begin{aligned}
\mathrm{Tr}\gamma_\mu\gamma_\nu\gamma_\alpha\gamma_\beta\gamma_\delta &= -2i(g_{\mu\nu}\epsilon_{\alpha\beta\delta} + g_{\alpha\beta}\epsilon_{\mu\nu\delta} + g_{\beta\delta}\epsilon_{\mu\nu\alpha} - g_{\delta\alpha}\epsilon_{\mu\nu\beta}) \\
\mathrm{Tr}\gamma_\mu\gamma_\nu\gamma_\alpha\gamma_\beta\gamma_\rho\gamma_\sigma &= -2(g_{\mu\nu}g_{\alpha\beta}g_{\rho\sigma} - g_{\mu\alpha}g_{\nu\beta}g_{\rho\sigma} + g_{\mu\beta}g_{\nu\alpha}g_{\rho\sigma} - g_{\alpha\beta}g_{\mu\rho}g_{\nu\sigma} \\
&\quad + g_{\alpha\beta}g_{\mu\sigma}g_{\nu\rho} - g_{\mu\nu}g_{\alpha\rho}g_{\beta\sigma} + g_{\mu\nu}g_{\alpha\sigma}g_{\beta\rho} - \epsilon_{\mu\nu\rho}\epsilon_{\alpha\beta\sigma} + \epsilon_{\mu\nu\sigma}\epsilon_{\alpha\beta\rho})
\end{aligned}
$$

and so on. The trace of an odd number of γ matrices does not vanish as it does in 3+1 dimensions.

Appendix B. Some useful relations

First, for the non-relativistic case

$$
\begin{aligned}
[\hat{\mathbf{\Pi}}_2, \frac{1}{H_0}] &= \frac{i}{ml^2}\frac{1}{H_0}\hat{\mathbf{\Pi}}_1\frac{1}{H_0} \\
[\hat{\mathbf{\Pi}}_1, \frac{1}{H_0}] &= -\frac{i}{ml^2}\frac{1}{H_0}\hat{\mathbf{\Pi}}_2\frac{1}{H_0} \\
[\hat{\tau}, \frac{1}{H_0}] &= \frac{1}{H_0^2}
\end{aligned}
\tag{70}
$$

For the relativistic case

$$
\begin{aligned}
[\hat{x}_0, Q_1] &= -4iS\frac{1}{M_0}S\hat{\Pi}_0\frac{1}{M_0} + 2i\hat{\Pi}_0\frac{1}{M_0} \\
[\hat{x}_0, Q_2] &= -\frac{4i}{l^2}\frac{1}{M_0}S\hat{\Pi}_0\frac{1}{M_0} \\
[\hat{\mathbf{\Pi}}_1, Q_2] &= -\frac{2i}{l^4} frac1M_0\{S, \hat{\mathbf{\Pi}}_2\}\frac{1}{M_0} \\
[\hat{\mathbf{\Pi}}_1, Q_1] &= -\frac{2i}{l^2}\frac{S}{M_0}\{S, \hat{\mathbf{\Pi}}_2\}\frac{1}{M_0} + \frac{2i}{l^2}\hat{\mathbf{\Pi}}_2\frac{1}{M_0} \\
[\hat{\mathbf{\Pi}}_2, Q_1] &= \frac{2i}{l^2}\left[\frac{1}{M_0}\{S, \hat{\mathbf{\Pi}}_1\}\frac{1}{M_0} - \hat{\mathbf{\Pi}}_1\frac{1}{M_0}\right] \\
[\hat{\mathbf{\Pi}}_2, Q_2] &= \frac{2i}{l^4}\frac{1}{M_0}\{S, \hat{\mathbf{\Pi}}_1\}\frac{1}{M_0}
\end{aligned}
\tag{71}
$$

References

[1] S. L. Shapiro and S. A. Teukolsky, "*Black Holes, White Dwarfs and Neutron Stars, The Physics of Compact Objects*", (Wiley, N.Y.) (1983).

[2] R. Narayan, P. Paczynski and T. Piran, *Ap. J. Lett.*, 395, **L**83 (1992).

[3] J. Ambjorn and P. Olesen, *Nucl. Phys.* **B** 315, 606, (1989), and references therein.

[4] V. V. Skalozub, *Sov. J. Nucl. Phys.* **45**, 1058, (1987) and referenc es therein.

[5] A. Erdas and G. Feldman, *Nucl. Phys.* **B** 343, (1990), 59 7.

[6] P.H. Damgaard and D. Espriu, *Phys. Lett.* **B** 246, 442, (1991).

[7] I.V. Krive and S. A. Naftulin, *Phys. rev.* **D** 46, 2737, (1992).

[8] K.G. Klimenko, *Z. Phys.* **C** 54, (1992), 323.

[9] P. K. Panigrahi, R. Ray and B. Sakita, *Phys. Rev.* **B** 42, 4036, (1990).

[10] A.M.J. Schakel, "On Broken Symmetries in Fermi Systems", (1989) and references therein.

[11] A. Neagu and A.M.J. Schakel, *Phys. Rev.* **D** 48, 1785, (1993).

[12] S. Iso, D. Karabali and B. Sakita *Phys. Lett.* **B** 296, 143, (1992); *Nuc. Phys.* **B** 388, 700 , (1992)

[13] R. Ray and G. Gat, work in progress.¸

[14] S. Sakhi, "*Effective action of a 2+1 dimensional system of nonrelativistic fermions in the presence of a uniform magnetic field: dissipation effects*" UdeM-LPN-TH-179.

[15] S. Deser, R. Jackiw and S. Templeton *Ann. Phys.* 140, 372, (1982).